



# INTERNATIONAL JOURNAL OF PHARMACEUTICAL RESEARCH AND NOVEL SCIENCES

# IJPRNS

## COST ANALYSIS OF EMERGENCY VISITS DUE TO DRUG RELATED PROBLEMS

Malini. S<sup>1</sup>, Pasupathi. A<sup>2</sup>

<sup>1</sup>Research Scholar, Vinayaka Mission Research Foundation  
(Deemed to be University), Salem, Tamil Nadu, India

<sup>2</sup>Department of Pharmaceutics, Vinayaka Mission college of Pharmacy, Salem, Tamil Nadu, India

### ABSTRACT

To identify patients coming to Emergency Medicine Department (EMD) with drug related problems, classify the DRPs and calculate the direct cost spent for treating them. This was a prospective observational study conducted in emergency medicine department. The patients coming to EMD with DRPs were classified according to Cipolle's classification and the direct medical and non medical costs were calculated. A total of around 107 patients identified with DRPs of which 99 patients were included in the study. In this study, 51% of the cases were due to ADR and 35% due to non adherence and rest of the cases were due to overdose (10%), drug interaction (3%) and sub therapeutic dose (1%). Major portion for treatment was spent for direct medical cost in which cost for laboratory investigations have contributed the most, INR 10, 93,992 (42%) followed by Health care professional cost INR 55, 6814 (21%), Pharmacy cost INR 4, 00,524.6 (15%), Admission cost INR 3, 80,400 (15%). The direct non medical cost includes cost for diet and travel which was found to be INR 1, 68,443 and INR 71,947 respectively. The drug related problems adds a significant economic burden on the patients which can be reduced by imparting knowledge about the proper use of medicines and by improving collaborative efforts of the patients, physicians, pharmacists and caregivers.

**Key Words:** Drug related problems (DRPs), Cost analysis, direct cost.

### Author for correspondence

**Malini. S,**

Research Scholar, Vinayaka Mission Research Foundation (Deemed to be University), Salem, Tamil Nadu, India

### INTRODUCTION

Medications are used in various disease conditions to optimize drug therapy with minimum safety related problems within the frame work of pharmaceutical care plan. Over the last few decades advances in drug therapies have improved patient care and there has been an apparent increase in the incidence of Drug Related Problems (DRPs) being reported [1]. DRP can

the health outcomes in the patients. It can occur during any stages of drug use starting from prescribing to administration. Lack of follow-ups may also leads to drug related problems [2]. Various risk factors for DRPs include polypharmacy, co morbid diseases, increased medicine use. Another major contributing factor is the practice of self medication with over the counter (OTC) medications available in India and many other parts of the world [3-5]. Even with the combined efforts taken by the manufacturers, regulatory authorities and health care professionals to reduce the incidence of DRPs, they continue to occur due to various reasons that can be attributed to different stages of drug use. DRPs occur in patients

because of the improper use that might appear at any level of drug use. This demands appropriate attention and interventions to reverse the condition which might expose a patient to severe harm than the disease itself for which one was taking the medicine. The severity of DRPs can range from minor side effects to death of the patient. Nowadays a lot of hospital visits and admissions can be attributed to DRPs and this rate seems to be rising over the past two decades. In the process of reversing the DRPs healthcare resources and a significant amount of money is spent which poses an economic burden on the patients as they have to spend more than what was intended for their ailment. The economic impact of DRPs is extensive on patients, healthcare providers, administrators and society. Estimates have shown that for every US \$1 spent on drugs, US \$1.33 is consumed to treat DRPs [6]. The cost of DRPs has increased every year and in 2008 it was estimated at US \$289 billion in the United States [7]. An Indian study showed total cost for the management of DRPs was calculated to be INR17,37,339. The direct cost was INR 1,72,961 and the approximate indirect cost was INR15,64, 378. The drugs used for treatment (54%) contribute significant part in direct cost. Among various drug related problems medication non adherence was the main contributor (85% of total cost) followed by adverse drug reactions (13% of total cost) [8]. In a developing country like ours, majority of the population belongs to middle class with only sufficient money for their day to day living and do not have medical insurance coverage. Also most of the medications are available as OTC drugs and lack of qualified pharmacy professionals for handling the medicine. Therefore, creating a clear picture of hospital visits due to drug related problems and its economic impact is

significant. Even though a number of studies have evaluated the pharmaco-economic impact of DRPs in the developed world, only limited data are available from developing countries. In our study, we have attempted to calculate the direct cost involved to treat patients who present to the Emergency department because of DRPs.

## **MATERIALS AND METHODS**

A prospective observational study was conducted over a period of six months in the Emergency medicine, Medical Centre, Kottayam, Kerala. The study was approved by the Institution Human Ethics Committee of the hospital. Patients came to emergency medicine department with drug related problems were selected for the study. Both genders and all age groups were included. Critically ill patients, patients who are not willing to participate and patient's with claim or insurance were excluded from the study.

Patient data collection form included demographic details, medical and medication history, direct medical and non-medical cost. The DRPs identified and were categorized based on CIPOLLEs classification, which include Adverse drug reactions(ADR), Medication error, Non compliance, Drug interactions, Untreated indication, Drug use without indication, Improper drug selection, Sub therapeutic Dose, Supra therapeutic Dose. The ADR's were classified based on the WHO-UMC scale and preventability of ADR was categorized based on the Schumock and Thornton scale. The direct cost for each DRPs were calculated. Pharmacy cost, Laboratory cost, Admission cost, Health care professional cost were included in the direct medical cost whereas direct non medical cost included Diet and Transport charges.

## **RESULTS AND DISCUSSION**

During our study period of 6 months, 10,485 patients visited the emergency department, of which 107 were due to drug related problems. From these cases, 8 were excluded after meeting the exclusion criteria and the remaining 99 cases were included in our study. The DRPs were found to be almost equally predominant in both genders, of which 48% were females and 52% were males. The 99 subjects enrolled in our study were classified into 5 age groups ranging from 0-18, 18-29, 40-59, 59-79 and 80-99 years. Most frequent occurrence of DRPs was observed in the age group 40-59, which was about 45%. The DRPs observed were grouped according to CIPOLLE's classification into 5 categories. The patients included in our study experienced different type of drug related problems like adverse drug reactions, non compliance or medication error, drug interactions, sub-therapeutic dose and over dose. In this

study, 50 patients presented with an ADR (51%), followed by 35 who were non-adherent to the medicines and medical advice provided (35%).Overdose of medications was observed in 10 patients (10%), while in 3 patients drug interactions was seen (3%) and 1 of the cases received a sub- therapeutic dose(1%) which is shown in figure-1.




Figure-1 Distribution Of DRP's as per Cipolle's Classification

In the DRPs, ADR showed the highest prevalence and was seen in 50 patients (51%). The ADR's were classified based on the WHO-UMC scale and it showed 64% was probable, 32% possible and 4% certain from the total number of ADR's. The preventability of ADR was categorized based on the Schumock and Thornton scale, which revealed 34% of ADR's were not preventable and 66% were preventable of which 30% were definitely preventable which is shown in the figure-2&3. This may be due to selection of drug, dose, route of administration, frequency, drug interaction, lack of patient's knowledge.




Figure-2 Preventability Of ADR




Figure-3 Preventability Class

Of the 99 cases assessed, 15 subjects were admitted due to DRPs associated with anti-diabetics (14%) followed by 14 admissions by anti-epileptics (13%) and anti-psychotics (13%) each. Direct cost spent in treating DRPs is an addition of direct medical cost and direct non medical cost. Direct medical cost includes pharmacy cost, healthcare professionals cost, laboratory cost, admission cost that contributed to almost 92% of the total and direct non-medical cost included diet and travel charges. A total direct cost of about INR 25, 96,007 was spent to treat the patients who visit the EMD due to drug related problems. .

The patients were treated in both inpatient (IP) and outpatient (OP) settings. For managing the DRPs for outpatients, INR 75,248 (33% of total direct cost) was spent and for those treated as inpatients, an amount of INR 25,20,829 (67% of total direct cost) was disbursed. A total of INR 23, 56,728 was spent for direct medical cost of which, 42% comprised of laboratory charges amounting to INR 10, 39,992. A direct non medical cost of INR 2,39,349 was spent by the patients during our study period.6% of the total direct cost was the amount spend towards diet which is the highest contributor in direct non- medical cost of INR 1,68,443 were shown in table-1 .

**Table-1 DISTRIBUTION OF DRPs IN TREATMENT COST**

Type of cost	Total cost (in INR)	Percentage (%)	Mean	Standard deviation (±)
Direct medical cost				
Pharmacy cost	400524.6	15%	4045.70	17849.91
Health care professionals cost	556814	21%	5624.38	14280.99
Laboratory cost	1093992	42%	11050.42	19525.63
Admission cost	380400	14%	3842.42	11268.86
TOTAL	2356728.148	92%	23805.33	47071.31
Direct non-medical cost				
Diet	168443	6%	1701.44	2319.83
Travel	71947	2%	726.73	768.90
Total	239349.2	8%	2417.66	2797.86
Grand total	2596077	100%	26223.00	48568.96

In the total direct cost, 67% was contributed by 66 in- patients and 33% by the out-patients. Even though direct cost for inpatients and out-patients was found to be similar, direct non medical cost (25%) was found to be a major contributing factor for out-patients. whereas direct medical cost was substantial for in- patients. In our study the direct cost spent for ADR was INR 15,20,960 (58%), which shows ADR is the highest contributor among the DRPs, then comes non adherence with INR 8,32,656 (32%) and for drug- interaction and overdose it was INR 1,93,888 (6%) and INR 1,23,052 (4%) respectively. The direct cost spent for ADR was about INR 15,20,960 , in which direct medical cost was INR 13,94,037 and direct non- medical cost was INR 1,26,923. Among various drug related problem's adverse drug reactions (ADR) was the main contributor (51%) of total cost, it was observed that the potential for an adverse reaction like cutaneous reaction like rash, generalized pruritus, was 43% followed by hepatotoxicity. ATT had the highest potential (44%) for an adverse drug reaction followed by steroid (42%). Non adherence to anti-diabetic, anti- epileptics, anti-tubercular, anti-hypertensive drugs were predominant DRPs reported, which escalates the treatment cost especially the direct medical cost. Approximately about INR 23790.17 was spent by a patient, in which direct medical cost was INR 21501.95 and direct non medical cost was INR 2288.22. The direct cost spent for drug- interaction was about INR 1,93,888, in which direct medical cost was INR 1,77,864 and direct non-medical cost was INR 16,024. The laboratory cost shows highest percentage was about 53%

from the total medical cost. Management of drug interaction leading to hospitalization incurred the least expenditure was about 3%.

Among the 10 overdose cases, major portion of the total cost was spent for healthcare professionals (34%), which mainly includes the fees for physicians, nurses, and physiotherapist and is about INR 4238. The direct cost spent for overdose was about INR 1,23,0522 in which direct medical cost was INR 1,11,60 and direct non-medical cost was INR 11,4480. The direct cost spent for sub-therapeutic dose was about INR 7972.49 in which direct medical cost was INR 7272.49 and direct non medical cost was INR 700.

The prevalence of DRPs was higher in males (52%) than in females (48%) which were also comparable to the population in an Indian study [8]. About 45% of DRP was observed in the age group 40-59 which may be due to poor knowledge about the disease and the drugs and other risk factors such as poly-pharmacy, co-morbid conditions and lack of follow-up and reassessment of medical treatment. Similarly, in a study by Baena MI *et al*, the patients within the age group of 45-64 taking 3 or more medicines presented an odds ratio of 64.07 [9]. Among the various DRPs, ADRs showed the highest prevalence. A study by Peter J Zed *et al* showed drug-related visits were predominantly due to adverse drug reactions, followed by non-adherence which was almost identical to our findings [10]; while overprescribing of the correct medication was found to be the major cause in the study by Tafreshi *et al* [11]. The distribution of ADRs based on preventability showed that most of the ADRs were preventable. Adverse drug related events accounted for 12% of emergency department visits, of which 68% were considered preventable in the first study [10]. Apart from these, a study from Karnataka showed that 47.55% of DRPs are due to drug interaction [12]. In our study most common drug classes that caused DRPs were anti-diabetics, anti- psychotics, anti-epileptics .Peter J Zed *et al*, in one of his study identified antimicrobial agents (11.2%), opioid containing analgesics (11.2%), antipsychotics (9.5%) and benzodiazepines (6.1%) as the major contributors of DRPs [10]. In his other study in 2018 the most common drug classes were anti-infective (27.4%), respiratory agents (22.4%), central nervous

system agents (20.4%), immunosuppressant's (7.5%), and gastrointestinal agents (6.0%) [13]. A study by Tafreshi revealed the most common medications involved were cardiovascular agents, narcotics, antibiotics, asthma medications, antidepressants, anti- epileptics, acetaminophen, anti-Parkinson medications[11]. In treating DRPs a significant amount of money is spent and it is actually an additional expense to the normal cost of drug use by the patients A study conducted in India by Smita Pattanaik *et al*, showed that the total cost of management of all the 92 drug related events was calculated to be INR 17,37,339. The major share of the direct cost was due to the drugs used for treatment of DRPs (54%). In addition, medication noncompliance was the single most important contributor to the cost (85% of the total cost), followed by ADRs (13% of the total cost), while management of drug overdose leading to hospitalization incurred the least expenditure [8]. When a person visits the hospital due to a DRP, if treated in inpatient setting, an average of about INR 38,194.38 is spent and if treated as outpatient, an average of about INR 2,280.243 is spent. The patients, based on their DRP's severity and the amount of attention needed to reverse the condition, they are decided whether they have to be treated in the outpatient setting or whether they have to be treated in the inpatient setting and accordingly the cost for the reversal of DRPs vary as mentioned above. Usually when the patient is severely affected, he/she might have to be admitted in the hospital in the care of various healthcare professionals to reverse the DRP or if the DRP is not that severe, like in allergies caused by the drugs, the patient is stabilized and then discharged without admission in the hospital.

## CONCLUSION

A large portion of DRPs observed were preventable cases of ADRs and non adherence to treatment. The most common drug classes involved were anti-diabetics, anti-epileptics, anti-psychotics and anti-hypertensive drugs which are used for the predominant diseases in India. Laboratory cost was

found to be the major contributor for direct medical cost. Educating patients about the correct use of their medications, showing the importance of adherence to treatment could reduce pressures on emergency services, hospitalizations, and consequently costs to the hospital, the health system, and the society.

## REFERENCE

1. M. Sonal Sekhar , C. Adheena Mary , P.G. Anju , Nishana Ameer Hamsa. Study on drug related hospital admissions in a tertiary care hospital in South India. *Saudi Pharmaceutical Journal* 2011; 19:273–8.
2. Adusumilli P K, Adepu R. drug related problems: an over view of various classification systems. *Asian Journal of Pharmaceutical and Clinical Research* 2014;7:7-10.
3. Sundus Basharat. Management of Drug Related Hospital Admissions. *International Journal of Scientific and Research Publications* 2016;6: 63- 7.
4. Shankar PR, Partha P, Shenoy N. Self-medication and non-doctor prescription practices in Pokhara valley, Western Nepal: A questionnaire-based study. *BMC Fam Pract* 2002; 3:17.
5. Selvaraj K, Kumar SG, Ramalingam A. Prevalence of self-medication practices and its associated factors in Urban Puducherry, India. *Perspect Clin Res.* 2014;5:32–6.
6. Johnson JA, Bootman JL. Drug-related morbidity and mortality: a cost-of-illness model. *Arch Intern Med* 1995; 155:1949– 56.
7. Gabriel Rodrigues Martins de Freitas, Mariana Younes Tramontina, Giacomo Balbinotto, Dyfrig Arwyn Hughes, Isabela Heineck, , Economic impact of Emergency Visits due to Drug-Related Morbidity on a Brazilian Hospital . *Value in Health Regional Issue* 2017;14C:1-8.
8. Smita Pattanaik, Punit Dhamija, Samir Malhotra, Navneet Sharma , Promila Pandhi . Evaluation of cost of treatment of drug related events in a public care tertiary sector. *Br J Clin Pharmacol* 2009;67:363-9.
9. Baena MI, Faus MJ, Fajardo PC, Luque FM, Sierra F, Martinez-Olmos J, Cabrera A, Fernandez-Llimos F, Martinez-Martinez F, Jiménez J, Zarzuelo A. Medicine-related problems resulting in emergency department visits. *Europe Journal Clinical Pharmacology* 2006; 62:387-93.
10. Peter J. Zed, Riyad B. Abu-Laban , Robert M. Balen , Peter S. Loewen , Corinne M. Hohl MD, Jeffrey R. Brubacher , Kerry Wilbur, Matthew O. Wiens , Leslie J. Samoy, Katie Lacaria , Roy A. Pursell. Incidence, severity and preventability of medication-related visits to the emergency department: a prospective study. *Canadian Medical Association or its licensors* 2008;178:1563-9.
11. Tafreshi MJ, Melby MJ, Kaback KR, Nord TC, Medication-Related Visits to the Emergency Department: A Prospective Study. *The Annals of Pharmacotherapy* 1999; 33:1252-7.,
12. Sarfaraz Mohammed. Sanjay Poudel, Fernance Laloo, Arjun madhur, Rinson Robert, Binu Mathew. Assessment of drug-related problems in a tertiary care teaching hospital, india. *Asian Journal of Pharmaceutical and Clinical Research* 2017; 10:310-13.
13. Peter J. Zed, Karen J. L. Black, Eleanor A. Fitzpatrick, Stacy Ackroyd-Stolarz, Nancy G. Murphy, Janet A. Curran, Neil J. MacKinnon, Doug Sinclair. Medication-Related Emergency Department Visits in Pediatrics: a Prospective Observational Study. *Official journals of the American academy of pediatrics* 2018; 135:435- 43.